

**BASF**

We create chemistry

# Juventus®

A fungicide for the control of a wide range of diseases in winter and spring wheat, durum wheat, winter and spring barley, triticale, rye, oilseed rape, peas, field beans and lupins; and a spring plant growth regulator in oilseed rape.

An emulsifiable concentrate containing 90 g/litre metconazole.

## Risk and Safety Information

### Warning

**Causes serious eye irritation.**  
**Suspected of damaging the unborn child.**  
**Harmful to aquatic life with long lasting effects.**

Obtain special instructions before use.  
Wear protective gloves/clothing/eye protection.  
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

IF exposed or concerned: Call a POISON CENTRE or doctor/physician.  
Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

PCS No. : 05780



® = Registered trademark of BASF

UFI : Y358-80JD-C004-W785

Supplied by:  
BASF Ireland DAC, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport, SK1 3GG, UK  
Telephone: 00 44 (0)161 475 3000  
Emergency Information (24 hours freephone): 0049 180 227 3112  
Technical Enquiries: 0044 (0)845 602 2553 (office hours)

Authorisation holder:  
BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport, SK1 3GG, UK Telephone: 00 44 (0)161 475 3000

# 5 L

81174346 IE 2024



**FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE and PLANT GROWTH REGULATOR, as directed below:**

<b>Crops</b>	<b>Maximum Individual Dose</b>	<b>Maximum Total Dose Per Crop</b>	<b>Latest Time of Application</b>
Winter wheat Spring wheat Durum wheat Winter barley Spring barley Triticale Rye	1.0 litre product/ha	2.0 litres product/ha	Up to and including GS 71 (caryopsis watery ripe stage)
Oilseed rape	0.8 litres product/ha	1.6 litres product/ha	10% of pods at final size
Combining peas Vining peas Field beans Lupins	0.8 litres product/ha	1.6 litres product/ha	14 days before harvest

**Other specific restrictions.**

A minimum interval of 14 days must be observed between applications on oilseed rape, peas, beans and lupin. A minimum interval of 21 days must be observed between applications on cereals.

To protect birds, only one application is allowed on cereals before GS 29 (end of tillering).

**READ ALL PRECAUTIONS BEFORE USE.**

**PRECAUTIONS**

**Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

**WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD)** when handling the concentrate.

**WEAR SUITABLE PROTECTIVE GLOVES** when handling contaminated surfaces.

**IF YOU FEEL UNWELL,** seek medical advice immediately (show the label where possible).

**WASH ALL PROTECTIVE CLOTHING** thoroughly after use, especially the insides of gloves.

**WHEN USING DO NOT EAT, DRINK OR SMOKE**

**Environmental protection**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies. Direct spray away from water.

**STORAGE**

**KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.**

**KEEP OUT OF REACH OF CHILDREN.**

**KEEP IN ORIGINAL CONTAINER,** tightly closed, in a safe place.

On emptying the container, **RINSE CONTAINER THOROUGHLY** by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

**DO NOT RE-USE CONTAINER FOR ANY OTHER PURPOSE.**

**PROTECT FROM FROST**

**DO NOT STORE IN DIRECT SUNLIGHT.**

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Juventus is a broad spectrum fungicide with curative and protectant activity, which can be used on all commercial varieties of winter and spring wheat, durum wheat, winter and spring barley, triticale and rye. Juventus can also be used on oilseed rape, field beans, combining and vining peas and lupins. When applied to oilseed rape in the spring, Juventus can provide plant growth regulation in the form of a reduction in crop height.

## Restrictions/Warnings

For professional use only.

For use by tractor mounted/trailed sprayers only

Do not apply to oilseed rape crops that are damaged or stressed from factors such as previous product use, adverse weather conditions (e.g. drought, frost or waterlogging), nutrient deficiency or pest attacks.

Juventus is formulated specifically for use on oilseed rape and legumes. The addition of further adjuvants is not advised.

Juventus should not be applied with pyrethroids to flowering oilseed rape.

Ensure the sprayer is free from previous chemical residues that may harm the crop. Use of a detergent cleaner is advised before and after use.

Avoid spray drift on to neighbouring crops and field margins.

## Resistance Management

GROUP	3	FUNGICIDE
-------	---	-----------

Juventus contains metconazole, a DMI fungicide. Resistance to some DMI fungicides has been identified in Septoria leaf blotch (*Mycosphaerella graminicola*) which may seriously affect the performance of some products. For further advice on resistance management in DMIs contact Teagasc, your agronomist or specialist advisor, and visit the FRAG-UK website.

Cereal Powdery mildew strains with decreased sensitivity to triazoles are common. Where these occur or develop, Juventus may not give satisfactory control.

For crop/pathogen situations where repeated spray applications are required, alternation or mixtures with an effective non cross-resistant fungicide are recommended. Fungicide

use does not replace the need for non-chemical cultural control methods e.g. resistant crop varieties, good agronomic practice, plant hygiene.

## Crop Specific Information

### Cereals

Juventus is a broad spectrum fungicide with curative and protectant activity which can be used on all commercial varieties of winter and spring wheat, durum wheat, winter and spring barley, triticale and rye for disease control as summarised below:

	Wheat	Barley	Triticale	Rye
Septoria leaf blotch ( <i>Mycosphaerella graminicola</i> )	MC	--	MC	--
Glume blotch ( <i>Septoria nodorum</i> )	MC	--	MC	--
Yellow rust	C	C	C	C
Brown rust	C	C	C	C
<i>Rhynchosporium secalis</i> (leaf scald)	--	R	--	R
Net blotch	--	R	--	--
Powdery mildew ( <i>Blumeria graminis</i> )	MC	MC	MC	MC
<i>Fusarium</i> ear blight	R	--	R	--

C = full control

MC = moderate control

R = reduction

### Time of Application

Apply as soon as diseases start to develop in the crop and repeat as required.

Application at an early stage of mildew development (not more than 3% infection on any green leaf) will give moderate control. Where mildew has become established, the use of a specific mildewicide will improve control and help prevent the development of resistant strains.

For optimum results on yellow rust apply before any leaf has more than 1% infection or as a preventative treatment on susceptible varieties after the flag leaf has fully emerged (GS 39).

Good reduction of ear Fusarium may be achieved when applied between full ear emergence and anthesis complete (GS 69).

For control of brown rust treat susceptible varieties as soon as rust is seen and spray more resistant varieties before any of the top three leaves have more than 1-2% infection.

Spray as soon as net blotch is seen on any of the top three leaves. A further treatment will be necessary where disease is well established

### Rate of Application

1.0 l/ha

## Oilseed rape

All varieties of oilseed rape may be treated.

### Disease control

Juventus provides control of *Alternaria*, and a reduction of phoma leaf spot, phoma stem canker (*Phoma lingam*) and light leaf spot.

	<i>Alternaria spp</i>	Phoma leaf spot & phoma stem canker ( <i>Phoma lingam</i> )	Light leaf spot
Oilseed rape	Control	Reduction	Reduction

### Time of Application

Apply as soon as diseases start to develop, and repeat as required.

### Rate of Application

0.8 l/ha.

### Crop height reduction

When applied to oilseed rape in the spring, Juventus can provide plant growth regulation in the form of a reduction in crop height.

### Time of Application

Apply during stem extension in the spring, once the oilseed rape crop is actively growing (GS 31-51)

### Rate of Application

0.8 l/ha.

## Combining Peas, Vining Peas, Field Beans and Lupins

All varieties of peas and field beans may be treated.

Consult processor before using on vining peas grown for processing.

Safety to lupins has not been established. If possible, a small area of the lupin crop should be treated before treating the whole crop.

### Disease control

Juventus can be used for the reduction of *Ascochyta* and *Mycosphaerella* species, the reduction of *Botrytis* and the control of rust on peas.

	<i>Ascochyta</i>	<i>Mycosphaerella</i>	<i>Botrytis</i>	Rust
Combining peas	Reduction	Reduction	Reduction	Control
Vining peas				
Field beans	-	-	-	Control

### Qualified Minor Use Recommendations

Juventus can be used for the reduction of *Ascochyta*, the reduction of *Botrytis* and the control of rust on lupins

	<i>Ascochyta</i>	<i>Mycosphaerella</i>	<i>Botrytis</i>	Rust
Lupins	Reduction	-	Reduction	Control

### Time of Application

	<i>Ascochyta</i>	<i>Mycosphaerella</i>	<i>Botrytis</i>	Rust
Combining peas	Start of flowering*	Start of flowering*	Mid flowering*	First sign of rust pustules*
Vining peas				
Field beans	-	-	-	Start of petal fall*
Lupins	Start of petal fall*	-	Start of petal fall*	Start of petal fall*

\* repeat 3-4 weeks later, if required.

### Rate of Application

0.8 l/ha.

## Following Crops

After treating a cereal, oilseed rape or legume crop with Juventus, the crops listed below may be sown as a following crop:

Beans	Clover	Oilseed rape	Ryegrass
Cabbage	Lettuce	Onions	Sugar beet
Carrots	Linseed	Peas	Sunflower
Cereals	Maize	Potatoes	

The effects of Juventus on other crops has not been assessed.

## Mixing and Application

### Application

In cereals apply Juventus in 200 to 300 l/ha water

In oilseed rape and legumes apply Juventus in 200-400 l/ha water.

Apply as a MEDIUM spray, as defined by BCPC. In dense crops use the higher volume to achieve adequate penetration and good coverage of the crop

To ensure optimum spray coverage and minimize spray drift, adjust the spray boom to the appropriate height above the crop, in according to guidance provided by the sprayer and/or nozzle manufacture.

Sprayers should be thoroughly cleaned out immediately after use

### Qualified approval

Juventus may be applied in reduced water volumes as specified in the table below, although efficacy at this reduced volume has not been evaluated. Application of this product at reduced-volume is at user's risk with regard to biological efficacy.

Crop	Rate	Minimum water volume
Winter wheat Spring wheat Durum wheat Winter barley Spring barley Triticale Rye	1.0 litres per hectare	110 litres per hectare

Crop	Rate	Minimum water volume
Oilseed rape Peas Beans Lupins	0.8 litres per hectare	90 litres per hectare

### Mixing

Never prepare more spray solution than is required.

Three quarters fill the tank with water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of Juventus to the spray tank while re-circulating. Fill up the tank with water and continue agitation until spraying is completed.

On emptying the container, rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to spray tank at time of filling and dispose of container safely.

### Compatibility

For information on compatibilities contact your distributor, local BASF representative or the BASF Technical Hotline 0044 (0) 845 602 2553. Alternatively, visit our website at [www.agricentre.basf.ie](http://www.agricentre.basf.ie)

Juventus should not be applied with pyrethroids to flowering oilseed rape.

All tank mixes should be used immediately after mixing.

### The following does not form part of the product label under S.I. No. 159 of 2012

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them

to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of any applicable law.

### **Additional Product Safety Information**

The information on this label is based on the best available information including data from test results.

### **Safety Data Sheet**

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



[agricentre.basf.ie/Juventus-IE/MSDS](http://agricentre.basf.ie/Juventus-IE/MSDS)

Alternatively, contact your supplier.

# Juventus®

**A fungicide for the control of a wide range of diseases in winter and spring wheat, durum wheat, winter and spring barley, triticale, rye, oilseed rape, peas, field beans and lupins; and a spring plant growth regulator in oilseed rape.**

An emulsifiable concentrate containing 90 g/litre metconazole.

## Risk and Safety Information

### Warning

**Causes serious eye irritation.  
Suspected of damaging the unborn child.  
Harmful to aquatic life with long lasting effects.**

Obtain special instructions before use.  
Wear protective gloves/clothing/eye protection.  
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

IF exposed or concerned: Call a POISON CENTRE or doctor/physician.  
Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

PCS No. : 05780



® = Registered trademark of BASF

UFI : Y358-80JD-C004-W785

Supplied by:  
BASF Ireland DAC, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport, SK1 3GG, UK  
Telephone: 00 44 (0)161 475 3000

Emergency Information (24 hours freephone): 0049 180 227 3112

Technical Enquiries: 0044 (0)845 602 2553 (office hours)

Authorisation holder:  
BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport, SK1 3GG, UK Telephone: 00 44 (0)161 475 3000

**5 L**