

Storm[®] Secure

BASF

We create chemistry



The convenient single feed, fixable block bait for rat and mouse control

Contains 0.05g/kg (0.005% w/w) flocoumafen
For use by Professionals only.
Not for sale to the general public

3 kg

© = Registered Trademark of BASF

Bait Blocks | Especially suitable for fixing in place

- ✓ Helps eliminate rodent infestations
- ✓ *Delivers a lethal dose in a single feed



811664361E2013

Storm® Secure

Contains 0.05 g/kg (0.005% w/w)
flocoumafen

Storm Secure is a ready-to-use block bait containing flocoumafen (0.005% w/w) for use as a rodenticide by Professionals for the control of rats and mice indoors and outdoors (around buildings only) for the protection of public health, stored products and materials.

For use by Professionals only.

Professional users must present their professional number such as herd/flock number, HFR or HPR number at point of sale to purchase this product for use only as specified on this label.

Danger:

May damage the unborn child.

May cause damage to organs (blood) through prolonged or repeated exposure.

If medical advice is needed, have product container or label at hand.

Keep out of reach of children.

Obtain special instructions before use.

Wear protective gloves.

IF exposed or concerned: Call a POISON CENTRE or doctor/physician.

Dispose of contents to comply with local requirements.

To avoid risks to human health and the environment, comply with the instructions for use.



Not for sale to the general public.

This product should only be used in accordance with a code of best practice such as the CRRU Ireland Best Practice Requirements for Rodent Control and Safe Use of Rodenticides. Follow any additional instructions in that code of best practice.

This product contains a bittering agent and a dye.

Use Biocides Safely and Sustainably.

It is illegal to use this product for uses or in a manner other than that prescribed on this label.

*Rodent death will occur in 4-10 days after ingestion of a lethal dose which may take a single feed.

Authorisation number: IE/BPA 70545

Batch number: see packaging

Expiry date: 3 years after manufacture

Authorisation Holder:

BASF Ireland DAC

Asgard House

19-20 City Quay

Dublin, D02 K744

Ireland

Tel: 0044 (0)161 475 3000

Emergency information: 0049 180 2273 112

(24 hours freephone)

Production date: see packaging



© = Registered trademark of BASF

UFI: 8CC0-Y0VG-U00H-4W9G



3 kg - 81176992|E2074

Application Method

Ready-to-use bait to be applied in tamper-resistant bait stations.

Application Rate:

Species	Situation	Application rate for each bait point	Frequency
House mouse (<i>Mus musculus</i>)	In and around buildings.	20g (1 block). If more than one bait station is needed, the minimum distance between bait stations should be of 1-2 meters.	Refer to instructions for Use below.
Brown rat (<i>Rattus norvegicus</i>) Black or roof rat (<i>Rattus rattus</i>)	In and around buildings.	40-60 g (2-3 blocks) If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters.	

Instructions for Use

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste).

Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

The product should be placed in the immediate vicinity of places where rodent activity has previously occurred (e.g. rodent runs, nesting sites, holes, burrows etc.).

Where possible, bait stations must be fixed to the ground or other structures.

Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see 'Risk Mitigation Measures' for the information to be shown on the label).

Bait should be secured so that it cannot be dragged away from the bait station.

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided
Protect bait from atmospheric conditions (e.g. rain, snow, etc). Place the bait stations/ baiting points in areas not liable to flooding.

Baits must be placed to minimise the exposure to non-target species and children.

Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

Wear protective chemical resistant gloves during product handling phase (e.g. EN 374).

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.

If after a treatment period of 35 days baits continue to be consumed and no decline in rodent activity is observed, the likely cause must be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.

For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
Remove the remaining product at the end of treatment period.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing of them.

The bait stations should be visited for mice at least every 2 to 3 days at the beginning and for rats only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Do not apply this product directly in the burrows.

Remove the remaining bait or the bait stations at the end of the treatment period.

Using this product should eliminate rodents within 35 days. In case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), seek advice from the product supplier or call a pest control service.

Risk Mitigation Measures

Where possible, prior to the treatment inform any bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign in accordance with the code of best practice.

Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

The baits must be securely deposited in a way so as to minimise the risk of consumption by other animals or children. Where possible secure baits so that they cannot be dragged away.

To reduce the risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment in line with the recommendations provided by the relevant code of best practice.

Do not use baits containing anticoagulant active substances as permanent baits or for the prevention of rodent infestation or monitoring of rodent activities.

Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.

Bait stations must be labelled with the following information: "Do not move or open"; "Contains a rodenticide"; "Product name or authorisation number"; "Active substance(s)" and "In case of incident, call the National Poisons Information Centre on (01) 809 2166".

Hazardous to wildlife.

Resistance Management

Do not use in areas where resistance to the active substance is suspected.

Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.

Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.

Storage

Store in a dry, cool and well ventilated place away from products with an odour.

Keep the container closed and away from direct sunlight.

Store in places prevented from the access of children, birds, pets and farm animals.

Shelf life: 3 years. Product is not to be stored at temperatures >30 °C.

Disposal of uneaten bait and dead rodents.

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with EPA requirements for the disposal of hazardous waste.

Use of gloves is recommended.

Dispose of dead rodents in accordance with local requirements, using one of the following methods of disposal (in order of preference): via an on-site or on-farm small carcass incinerator; with the site or farm's domestic waste; in the site or farm's normal non-hazardous waste; or by burial on-site, but away from sensitive areas.

Do not reuse the container for any other purpose.

Poison Information

For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-809 2166), retain the label for reference.

First Aid

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

Antidote: Vitamin K1 administered by medical/veterinary personnel only.

In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Contact a veterinary surgeon in case of ingestion by a pet.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below.



agricentre.basf.ie/Storm-Secure-IE/MSDS

Alternatively, contact your supplier.

Storm® Secure

Contains 0.05 g/kg (0.005% w/w)
flocoumafen

Storm Secure is a ready-to-use block bait containing flocoumafen (0.005% w/w) for use as a rodenticide by Professionals for the control of rats and mice indoors and outdoors (around buildings only) for the protection of public health, stored products and materials.

For use by Professionals only.

Professional users must present their professional number such as herd/flock number, HFR or HPR number at point of sale to purchase this product for use only as specified on this label.

Danger:

May damage the unborn child.

May cause damage to organs (blood) through prolonged or repeated exposure.

If medical advice is needed, have product container or label at hand.

Keep out of reach of children.

Obtain special instructions before use.

Wear protective gloves.

IF exposed or concerned: Call a POISON CENTRE or doctor/physician.

Dispose of contents to comply with local requirements.

To avoid risks to human health and the environment, comply with the instructions for use.



Not for sale to the general public.

This product should only be used in accordance with a code of best practice such as the CRRU Ireland Best Practice Requirements for Rodent Control and Safe Use of Rodenticides. Follow any additional instructions in that code of best practice.

This product contains a bittering agent and a dye.

Use Biocides Safely and Sustainably.

It is illegal to use this product for uses or in a manner other than that prescribed on this label.

*Rodent death will occur in 4-10 days after ingestion of a lethal dose which may take a single feed.

Authorisation number: IE/BPA 70545

Batch number: see packaging

Expiry date: 3 years after manufacture

Authorisation Holder:

BASF Ireland DAC

Asgard House

19-20 City Quay

Dublin, D02 K744

Ireland

Tel: 0044 (0)161 475 3000

Emergency information: 0049 180 2273 112

(24 hours freephone)

Production date: see packaging



© = Registered trademark of BASF

UFI: 8CC0-Y0VG-U00H-4W9G

3 kg - 81176992|E2074